

Welcome to Teva Shared Solutions®





1-800-887-8100 | M-F, 8 AM to 8 PM CT

APPROVED USES

AUSTEDO® XR (deutetrabenazine) extended-release tablets and AUSTEDO® (deutetrabenazine) tablets are prescription medicines that are used to treat:

- the involuntary movements (chorea) of Huntington's disease. AUSTEDO XR and AUSTEDO do not cure the cause of the involuntary movements, and it does not treat other symptoms of Huntington's disease, such as problems with thinking or emotions.
- movements in the face, tongue, or other body parts that cannot be controlled (tardive dyskinesia).
 It is not known if AUSTEDO XR and AUSTEDO are safe and effective in children.

IMPORTANT SAFETY INFORMATION

AUSTEDO XR and AUSTEDO can cause serious side effects in people with Huntington's disease, including: depression, suicidal thoughts, or suicidal actions. Do not start taking AUSTEDO XR or AUSTEDO if you are depressed (have untreated depression or depression that is not well controlled by medicine) or have suicidal thoughts. Pay close attention to any changes, especially sudden changes, in mood, behaviors, thoughts or feelings. This is especially important when AUSTEDO XR or AUSTEDO is started and when the dose is changed. Call your healthcare provider right away if you become depressed, have unusual changes in mood or behavior, or have thoughts of suicide.

Please read **Important Safety Information** on pages 6-7 and click **here** or visit www.AUSTEDO.com to read or print the **Medication Guide** for AUSTEDO XR.




What Is Teva Shared Solutions®?

Shared Solutions helps patients who are prescribed AUSTEDO XR extended-release tablets or AUSTEDO® (deutetrabenazine) tablets get started and stay on therapy by providing help with navigating insurance (including Medicare Part D), financial assistance offerings, refill support, and dedicated nurse support. You'll get real help from real people.

How to Enroll:



Patients can self-enroll at **MySharedSolutions.com**, or the healthcare provider can enroll the patient by completing a Prescription and Service Request Form (PSRF) found at **AUSTEDOHCP.com** or digitally within the CoverMyMeds portal.





What to Expect After Enrollment

A welcome call will come from Teva **Shared Solutions***. Save the number for future use.

• Look for **Teva ShSol** displayed on the phone or either of these phone numbers:

1-913-601-6146



1-800-887-8100



If you were enrolled in an office, your healthcare provider might have indicated that you may be interested in additional services, such as:

- Help understanding your insurance benefits and identifying in-network pharmacies and support through the prior authorization process
- Identification of financial assistance offerings (Free Trial Voucher, copay card, Patient Assistance Program)*


^{*}For eligible patients.



Dedicated Nurse Support

All patients have access to nurse support. After starting the prescription, the patient will get a call from a Teva **Shared Solutions**® nurse who will provide ongoing support with things like:

- Education about the diagnosis and prescription
- Treatment adherence, treatment expectations, dosing, and titration support
- Tools and resources to help manage treatment



It can be overwhelming to figure out how a treatment works; that's why we set up a program to support our patients every step of the way.

-Teva Shared Solutions® nurse



APPROVED USES

AUSTEDO® XR (deutetrabenazine) extended-release tablets and AUSTEDO® (deutetrabenazine) tablets are prescription medicines that are used to treat:

- the involuntary movements (chorea) of Huntington's disease. AUSTEDO XR and AUSTEDO do not cure the cause of the involuntary movements, and it does not treat other symptoms of Huntington's disease, such as problems with thinking or emotions.
- movements in the face, tongue, or other body parts that cannot be controlled (tardive dyskinesia).

It is not known if AUSTEDO XR and AUSTEDO are safe and effective in children.

IMPORTANT SAFETY INFORMATION

AUSTEDO XR and AUSTEDO can cause serious side effects in people with Huntington's disease, including: depression, suicidal thoughts, or suicidal actions. Do not start taking AUSTEDO XR or AUSTEDO if you are depressed (have untreated depression or depression that is not well controlled by medicine) or have suicidal thoughts. Pay close attention to any changes, especially sudden changes, in mood, behaviors, thoughts or feelings. This is especially important when AUSTEDO XR or AUSTEDO is started and when the dose is changed. Call your healthcare provider right away if you become depressed, have unusual changes in mood or behavior, or have thoughts of suicide.

Do not take AUSTEDO XR or AUSTEDO if you:

- have Huntington's disease and are depressed or have thoughts of suicide.
- have liver problems.
- are taking reserpine. Do not take medicines that contain reserpine with AUSTEDO XR or AUSTEDO. If your healthcare provider plans to switch you from taking reserpine to AUSTEDO XR or AUSTEDO, you must wait at least 20 days after your last dose of reserpine before you start taking AUSTEDO XR or AUSTEDO.
- are taking a monoamine oxidase inhibitor (MAOI) medicine. Do not take an MAOI within 14 days after you stop taking AUSTEDO XR or AUSTEDO. Do not start AUSTEDO XR or AUSTEDO if you stopped taking an MAOI in the last 14 days. Ask your healthcare provider or pharmacist if you are not sure.
- are taking tetrabenazine. If your healthcare provider plans to switch you from tetrabenazine to AUSTEDO XR or AUSTEDO, take your first dose of AUSTEDO XR or AUSTEDO on the day after your last dose of tetrabenazine.
- are taking valbenazine.



IMPORTANT SAFETY INFORMATION (CONTINUED)

Other possible serious side effects include:

- Irregular heartbeat (QT prolongation). AUSTEDO XR and AUSTEDO® (deutetrabenazine) tablets increases your chance of having certain changes in the electrical activity in your heart. These changes can lead to a dangerous abnormal heartbeat. Taking AUSTEDO XR or AUSTEDO with certain medicines may increase this chance.
- Neuroleptic Malignant Syndrome. Call your healthcare provider right away and go to
 the nearest emergency room if you develop these signs and symptoms that do not
 have another obvious cause: high fever, stiff muscles, problems thinking, very fast or
 uneven heartbeat, or increased sweating.
- Restlessness. You may get a condition where you feel a strong urge to move. This is called akathisia.
- Parkinsonism. Symptoms include: slight shaking, body stiffness, trouble moving, trouble keeping your balance, or falls.

Sleepiness (sedation) is a common side effect of AUSTEDO XR and AUSTEDO. While taking AUSTEDO XR or AUSTEDO, do not drive a car or operate dangerous machinery until you know how AUSTEDO XR or AUSTEDO affects you. Drinking alcohol and taking other drugs that may also cause sleepiness while you are taking AUSTEDO XR or AUSTEDO may increase any sleepiness caused by AUSTEDO XR and AUSTEDO.

The most common side effects of AUSTEDO in people with Huntington's disease include sleepiness (sedation), diarrhea, tiredness, and dry mouth.

The most common side effects of AUSTEDO in people with tardive dyskinesia include inflammation of the nose and throat (nasopharyngitis) and problems sleeping (insomnia).


The most common side effects of AUSTEDO XR are expected to be similar to AUSTEDO in people with Huntington's disease or tardive dyskinesia.

These are not all the possible side effects of AUSTEDO XR or AUSTEDO. Call your doctor for medical advice about side effects. You are encouraged to report side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Please read the accompanying Medication Guide.



Sign up for Teva Shared Solutions®



For patients, visit

MySharedSolutions.com

or contact us by phone

Monday-Friday from 8 AM

to 8 PM CT at 1-800-887-8100.



For healthcare providers, access the Prescription and Service Request Form (PSRF) here, obtain the patient's signature, and then fax the form to 1-844-257-6126.

Shared Solutions enrollment is also available within the CoverMyMeds portal.

