

Chorea is one of the most visible symptoms of Huntington's disease (HD) and can be treated. An important step in managing your HD chorea is talking to your doctor about treatment.



Answer the questions below, and bring this with you to your next doctor's appointment.

## 1 Which of the following involuntary movements are you experiencing?

Select all that apply.

- |  |  |  |                                    |
|--|--|--|------------------------------------|
| <input type="checkbox"/> Flicking the fingers                | <input type="checkbox"/> Shrugging the shoulders   | <input type="checkbox"/> Jerking and/or twisting | <input type="checkbox"/> Grimacing |
| <input type="checkbox"/> Irregular eye movements or blinking | <input type="checkbox"/> Flailing arms and/or legs |  |                                    |

## 2 How often do you experience these involuntary movements?

Please select one:

- |  |  |   |                                |
|--|--|---|--------------------------------|
| <input type="checkbox"/> Multiple times per minute | <input type="checkbox"/> Multiple times per hour | <input type="checkbox"/> Multiple times per day | <input type="checkbox"/> Other |
|--|--|---|--------------------------------|

## 3 Do you have trouble with any of these daily activities?

Select all that apply.

- |   |  |   |                                |
|---|--|---|--------------------------------|
| <input type="checkbox"/> Getting dressed    | <input type="checkbox"/> Brushing your teeth | <input type="checkbox"/> Speaking clearly         | <input type="checkbox"/> Other |
| <input type="checkbox"/> Drinking beverages | <input type="checkbox"/> Bathing             | <input type="checkbox"/> Swallowing and/or eating | <input type="checkbox"/> None  |

## 4 Does someone help you with everyday tasks?

Please select one:

- |                              |                             |
|------------------------------|-----------------------------|
| <input type="checkbox"/> Yes | <input type="checkbox"/> No |
|------------------------------|-----------------------------|

## 5 Has a loved one or care partner noticed your unintentional movements?

Please select one:

- |                              |                             |
|------------------------------|-----------------------------|
| <input type="checkbox"/> Yes | <input type="checkbox"/> No |
|------------------------------|-----------------------------|

### APPROVED USE

AUSTEDO® XR (deutetrabenazine) extended-release tablets and AUSTEDO® (deutetrabenazine) tablets are prescription medicines that are used to treat the involuntary movements (chorea) of Huntington's disease. AUSTEDO XR and AUSTEDO do not cure the cause of the involuntary movements, and it does not treat other symptoms of Huntington's disease, such as problems with thinking or emotions.

It is not known if AUSTEDO XR and AUSTEDO are safe and effective in children.

### IMPORTANT SAFETY INFORMATION

**AUSTEDO XR and AUSTEDO can cause serious side effects in people with Huntington's disease, including: depression, suicidal thoughts, or suicidal actions. Do not start taking AUSTEDO XR or AUSTEDO if you are depressed (have untreated depression or depression that is not well controlled by medicine) or have suicidal thoughts.** Pay close attention to any changes, especially sudden changes, in mood, behaviors, thoughts or feelings. This is especially important when AUSTEDO XR or AUSTEDO is started and when the dose is changed. Call your healthcare provider right away if you become depressed, have unusual changes in mood or behavior, or have thoughts of suicide.

**AUSTEDO XR (deutetrabenazine) extended-release tablets is a once-daily treatment for adults with HD chorea**  
Use these questions to talk to your doctor and learn more about treating Huntington's disease (HD) chorea.

**1 Will my HD chorea get worse over time?**

---

---

**2 How may AUSTEDO XR help me?**

---

---

**3 What is the Total Maximal Chorea (TMC) score? How is it used to determine the effectiveness of HD chorea treatment?**

---

---

**4 How does the dosing schedule work with AUSTEDO XR, and how will it impact my HD chorea treatment?**

---

---

**5 How do I take AUSTEDO XR?**

---

---

**6 Can you tell me about possible side effects of AUSTEDO XR?**

---

---

**7 What patient and care partner support will be available during AUSTEDO XR treatment?**

---

---

## IMPORTANT SAFETY INFORMATION (CONTINUED)

**Do not take AUSTEDO XR or AUSTEDO® (deutetrabenazine) tablets if you:**

- have Huntington's disease and are depressed or have thoughts of suicide.
- have liver problems.
- are taking reserpine. **Do not** take medicines that contain reserpine with AUSTEDO XR or AUSTEDO. If your healthcare provider plans to switch you from taking reserpine to AUSTEDO XR or AUSTEDO, you must wait at least 20 days after your last dose of reserpine before you start taking AUSTEDO XR or AUSTEDO.
- are taking a monoamine oxidase inhibitor (MAOI) medicine. **Do not** take an MAOI within 14 days after you stop taking AUSTEDO XR or AUSTEDO. **Do not** start AUSTEDO XR or AUSTEDO if you stopped taking an MAOI in the last 14 days. Ask your healthcare provider or pharmacist if you are not sure.
- are taking tetrabenazine. If your healthcare provider plans to switch you from tetrabenazine to AUSTEDO XR or AUSTEDO, take your first dose of AUSTEDO XR or AUSTEDO on the day after your last dose of tetrabenazine.
- are taking valbenazine.

**Other possible serious side effects include:**

- **Irregular heartbeat (QT prolongation).** AUSTEDO XR and AUSTEDO increases your chance of having certain changes in the electrical activity in your heart. These changes can lead to a dangerous abnormal heartbeat. Taking AUSTEDO XR or AUSTEDO with certain medicines may increase this chance.
- **Neuroleptic Malignant Syndrome.** Call your healthcare provider right away and go to the nearest emergency room if you develop these

signs and symptoms that do not have another obvious cause: high fever, stiff muscles, problems thinking, very fast or uneven heartbeat, or increased sweating.

- **Restlessness.** You may get a condition where you feel a strong urge to move. This is called akathisia.
- **Parkinsonism.** Symptoms include: slight shaking, body stiffness, trouble moving, trouble keeping your balance, or falls.

**Sleepiness (sedation) is a common side effect of AUSTEDO XR and AUSTEDO.** While taking AUSTEDO XR or AUSTEDO, do not drive a car or operate dangerous machinery until you know how AUSTEDO XR or AUSTEDO affects you. Drinking alcohol and taking other drugs that may also cause sleepiness while you are taking AUSTEDO XR or AUSTEDO may increase any sleepiness caused by AUSTEDO XR and AUSTEDO.

**The most common side effects of AUSTEDO in people with Huntington's disease include** sleepiness (sedation), diarrhea, tiredness, and dry mouth.

**The most common side effects of AUSTEDO in people with tardive dyskinesia include** inflammation of the nose and throat (nasopharyngitis) and problems sleeping (insomnia).

The most common side effects of AUSTEDO XR are expected to be similar to AUSTEDO in people with Huntington's disease or tardive dyskinesia.

These are not all the possible side effects of AUSTEDO XR or AUSTEDO. Call your doctor for medical advice about side effects. You are encouraged to report side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch) or call **1-800-FDA-1088**.

Please click [here](#) or visit [www.AUSTEDO.com](http://www.AUSTEDO.com) to read or print the **Medication Guide** for AUSTEDO XR.